

EPR of Yb³⁺ ions in Ba_{1-x}La_xF_{2+x} mixed crystals

Aminov L., Abdulsabirov R., Gafurov M., Korableva S., Kurkin I., Kurzin S., Rakhmatullin R., Ziganshin A.

Kazan Federal University, 420008, Kremlevskaya 18, Kazan, Russia

Abstract

Electron paramagnetic resonance (EPR) spectra of impurity Yb³⁺ ions (about 0.1 at.%) in mixed crystals BaF₂(1 - x) plus LaF₃(x) have been investigated for different values of the concentration x at a frequency of about 9.5 GHz by both continuous-wave (CW) EPR and electron spin echo methods. A spectrum of trigonal symmetry with a complex hyperfine structure is observed in "pure" BaF₂:Yb³⁺ (x = 0). Upon admixture of small amounts of LaF₃ (x = 0.001), additional EPR lines arise with intensities increasing with the increase of x up to 0.005. These lines are attributed to trigonal centers including two rare-earth ions and two compensating fluorine ions. A further increase of x results in a decrease of the total EPR spectrum intensity, and at x ≥ 0.05 the CW resonance becomes practically unobservable. This may be due to the formation of rare-earth ion clusters with paramagnetic Yb³⁺ ions occurring in domains with a disordered structure of surroundings resulting in very broad EPR lines, which cannot be registered by CW EPR. Indeed, very broad (not less than 1 kG) EPR lines were observed by the electron spin echo method for concentrations x ≥ 0.02. © Springer-Verlag 2005.
